

# Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

CODE: 7220

NAME: Petrifying springs with tufa formation (Cratoneurion)

## 1. National Level

### 1.1 Maps

1.1.1 Distribution Map	Yes
1.1.2 Distribution Method	Estimate based on partial data with some extrapolation and/or modelling (2)
1.1.3 Year or period	2007-2012
1.1.4 Additional map	No
1.1.5 Range Map	Yes

## 2. Biogeographical Or Marine Level

### 2.1 Biogeographical Region

#### Pannonian (PAN)

### 2.2 Published

Bölöni J., Molnár Zs. & Kun A (2011): Magyarország Élőhelyei Vegetációtípusok leírása és határozója ÁNÉR 2011: MTA Ökológiai és Botanikai Kutatóintézete, Vácrátót.

Molnár, Zs., M. Biró, J. Bölöni & F. Horváth (2008): Distribution of the (semi-) natural habitats in Hungary I.: Marshes and grasslands, Acta Botanica Hungarica 50 (Suppl): 59-105.

A Nemzeti Biodiverzitás-monitorozó Rendszer keretében 2007-2012 között végzett felmérések kutatási jelentése

### 2.3 Range of the habitat type in the biogeographical region or marine region

2.3.1 Surface area - Range (km <sup>2</sup> )	2799
2.3.2 Range method used	Estimate based on partial data with some extrapolation and/or modelling (2)
2.3.3 Short-term trend period	2001-2012
2.3.4 Short-term trend direction	increase (+)
2.3.5 Short-term trend magnitude	min max
2.3.6 Long-term trend period	
2.3.7 Long-term trend direction	N/A
2.3.8 Long-term trend magnitude	min max
2.3.9 Favourable reference range	area (km <sup>2</sup> ) operator approximately equal to (≈) unkown No method
2.3.10 Reason for change	Genuine Improved knowledge/more accurate data

### 2.4 Area covered by Habitat

2.4.1 Surface area (km <sup>2</sup> )	0,3
2.4.2 Year or period	2007-2012
2.4.3 Method used	Estimate based on partial data with some extrapolation and/or modelling (2)
2.4.4 Short-term trend period	2001-2012
2.4.5 Short-term trend direction	increase (+)
2.4.6 Short-term trend magnitude	min max

# Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

2.4.7 Short term trend method used	Estimate based on partial data with some extrapolation and/or modelling (2)	
2.4.8 Long-term trend period		
2.4.9 Long-term trend direction	N/A	
2.4.10 Long-term trend magnitude	min	max
2.4.11 Long term trend method used	N/A	
2.4.12 Favourable reference area	area (km) operator more than (>) unknown No method	
2.4.13 Reason for change	Genuine Improved knowledge/more accurate data	

## 2.5 Main Pressures

Pressure	ranking	pollution qualifier(s)
Landfill, land reclamation and drying out, general (J02.01)	high importance (H)	N/A
Modification of hydrographic functioning, general (J02.05)	high importance (H)	N/A
Canalisation & water deviation (J02.03)	medium importance (M)	N/A
invasive non-native species (I01)	medium importance (M)	N/A
Changes in abiotic conditions (M01)	medium importance (M)	N/A
Outdoor sports and leisure activities, recreational activities (G01)	medium importance (M)	N/A
damage caused by game (excess population density) (F03.01.01)	medium importance (M)	N/A

2.5.1 Method used – pressures based exclusively or to a larger extent on real data from sites/occurrences or other

## 2.6 Main Threats

Threat	ranking	pollution qualifier(s)
Landfill, land reclamation and drying out, general (J02.01)	high importance (H)	N/A
Modification of hydrographic functioning, general (J02.05)	high importance (H)	N/A
Canalisation & water deviation (J02.03)	medium importance (M)	N/A
invasive non-native species (I01)	medium importance (M)	N/A
Changes in abiotic conditions (M01)	medium importance (M)	N/A
Outdoor sports and leisure activities, recreational activities (G01)	medium importance (M)	N/A
damage caused by game (excess population density) (F03.01.01)	medium importance (M)	N/A

2.6.1 Method used – threats expert opinion (1)

## 2.7 Complementary Information

### 2.7.1 Species

Carex lepidocarpa

Carex appropinquata

Carex paniculata

Crotoneuron spp.

Pinguicula vulgaris

# Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

Nasturtium officinale

Potamogeton coloratus

## 2.7.2 Species method used

NBmR 5×5 km-es kvadrátok és N2000 területek élőhelytérképezése, az NBmR monitorozásra kiválasztott társulásainak cönológiai felvételezése, valamint a közösségi jelentőségű élőhelytípusok monitorozása eredményeinek összegzése és értékelése alapján.

## 2.7.3 Justification of % - thresholds for trends

## 2.7.4 Structure and functions - methods used

Estimate based on partial data with some extrapolation and/or modelling (2)

## 2.7.5 Other relevant information

A struktúra-funkció megítélése 5 komponensű (fajkészlet, fragmentáltság, inváziós fertőzöttség, termőhelyi sérülékenység, kezelések sikeressége) szempontrendszer alapján történt.

## 2.8 Conclusions (assessment of conservation status at end of reporting period)

### 2.8.1 Range

assessment Favourable (FV)  
qualifiers N/A

### 2.8.2 Area

assessment Inadequate (U1)  
qualifiers improving (+)

### 2.8.3 Specific structures and functions (incl Species)

assessment Inadequate (U1)  
qualifiers improving (+)

### 2.8.4 Future prospects

assessment Inadequate (U1)  
qualifiers improving (+)

### 2.8.5 Overall assessment of Conservation Status

Inadequate (U1)

### 2.8.5 Overall trend in Conservation Status

improving (+)

## 3. Natura 2000 coverage conservation measures - Annex I habitat types on biogeographical level

### 3.1 Area covered by habitat

#### 3.1.1 Surface area (km<sup>2</sup>)

min 0,3 max 0,3

#### 3.1.2 Method used

Estimate based on partial data with some extrapolation and/or modelling (2)

#### 3.1.3. Trend of surface area

N/A

### 3.2 Conversation Measures

#### 3.2.1 Measure

#### 3.2.2 Type

#### 3.2.3 Ranking

#### 3.2.4 Location

#### 3.2.5 Broad Evaluation

Other wetland-related measures (4.0)

One-off

medium importance (M)

Both

Long term

Managing water abstraction (4.3)

Administrative One-off

high importance (H)

Inside

Maintain Long term

